
Msx1 and Msx2 function together in the regulation of primordial germ cell migration in the mouse.

Journal: Dev Biol

Publication Year: 2016

Authors: Jingjing Sun, Man-Chun Ting, Mamoru Ishii, Robert Maxson

PubMed link: 27435625

Funding Grants: CIRM Stem Cell Biology Training Grant

Public Summary:

Scientific Abstract:

Primordial germ cells (PGCs) are a highly migratory cell population that gives rise to eggs and sperm. Much is known about PGC specification, but less about the processes that control PGC migration. In this study, we document a deficiency in PGC development in embryos carrying global homozygous null mutations in *Msx1* and *Msx2*, both immediate downstream effectors of Bmp signaling pathway. We show that *Msx1*(-/-);*Msx2*(-/-) mutant embryos have defects in PGC migration as well as a reduced number of PGCs. These phenotypes are also evident in a *Mesp1*-Cre-mediated mesoderm-specific mutant line of *Msx1* and *Msx2*. Since PGCs are not marked in *Mesp1*-lineage tracing, our results suggest that *Msx1* and *Msx2* function cell non-autonomously in directing PGC migration. Consistent with this hypothesis, we noted an upregulation of fibronectin, well known as a mediator of cell migration, in tissues through which PGCs migrate. We also noted a reduction in the expression of *Wnt5a* and an increase in the expression in *Bmp4* in such tissues in *Msx1*(-/-);*Msx2*(-/-) mutants, both known effectors of PGC development.

Source URL: <https://www.cirm.ca.gov/about-cirm/publications/msx1-and-msx2-function-together-regulation-primordial-germ-cell-migration>